

Plant Biodiversity and Conservation of Forests in Foot Hills of Garhwal Himalaya

Plant Biodiversity and Conservation of Forests in Foot Hills of Garhwal Himalaya

Munesh Kumar and Vishwapati Bhatt

Department of Forestry, Mizoram University, Post Box-190, Mizoram Aizwal-796009

E-mail: muneshmzu@yahoo.com

Department of Botany, Plant Physiology laboratory HNB, Garhwal University, Srinagar Garhwal

email: bhattvp03@yahoo.com

December 2006

Download at: <http://www.lyonia.org/downloadPDF.php?pdfID=2.490.1>

Plant Biodiversity and Conservation of Forests in Foot Hills of Garhwal Himalaya

Abstract

Floristic diversity, dominance and abundance to frequency ratio of tree, sapling, seedling, shrub and herb species were studied in two different forest sites of a tropical foot hill region of Garhwal Himalaya. In tree layer on both the sites the dominant species recorded were *Lannea coromandelica* (IVI-39.80) and *Anogeissus latifolia* (IVI-29.50) on site I and site II respectively. The ranges of diversity for tree layers was 4.580 to 4.643. Most of the species on both the sites were contagiously distributed except few species which were distributed randomly.

Introduction

The temperate forests of western and central Himalaya are usually distributed between 1200 and 3000 m asl which is the preferred zone by habitation and characterized extensive oak and coniferous forests. Puri (1960) considered that these forests represent climatic climax of one or other species of *Quercus* in upper altitudinal zones. The lower elevations of the temperate forests are occupied by oak-pine mixed forests and *Quercus semicarpifolia* with other coniferous at higher altitudes, normally form the climax vegetation. Other species of oak are found above the oak-pine mixed forests of Garhwal Himalaya (Osmostan 1922). The oaks are the most preferred tree species in the entire region and used mainly for fuel, fodder and small timber. In the lower altitudes (upto 1000m) generally the forests are dominated by *Shorea robusta*, alongwith *Anogeissus latifolia*, *Terminalia species* and *Adina cordifolia* (Kumar et al. 2004).

The various changes in the Himalayan forests are appearing in their structure, density and composition due to global warming (Gaur 1982), uncontrolled lopping and felling of trees for fuel wood, fodder and grazing (Bargali et al. 1998; Kumar et al. 2004). These biotic pressures which play an important role in forest community dynamics (Whitemore 1984; Pickett and White 1985) often regulate the recruitment and survival pattern of tree seedlings (Canham and Marks 1985). Each form of biotic pressure has different effects on the subsequent development of vegetation (Loucks et al.1980; Pandey and Singh 1985).

Several workers reported the status of forests and their management in the Himalayan region (Singh et al. 1981; Ralhan et al. 1982; Saxena and Singh 1982; Saxena et al. 1984) however, these studies are very limited to population structure in the part of Garhwal Himalaya and other parts of Central Himalaya. Therefore, the present paper deals with the analysis of vegetation and biotic pressure on forest in Kalsi range of Chakarata Forest division of Garhwal Himalaya.

Materials and Methods

Location and climate

The present study was carried out in Kalsi range of Chakarata Forest division (Yamuna valley), in two different sites i.e., site-I and site-II for comparative study of community composition and plant diversity, which is the part of the undulating terrains of Garhwal Himalaya (latitude 30 degrees 31' N and longitude 76 degrees 56' E) in District Dehradun of the Uttaranchal State in India. Altitudinally the region stretches from 700 to 1200 m asl, the Jaunsar-Bawar region which is remote and mysterious virtually isolated from rest part of the world. Since time immemorial, this mountainous region in the northern part of Uttaranchal has nurtured a unique life style.

Climatically spring, summer, rainy and winter seasons are well marked in this region. The maximum average rain fall is experienced during July and August. The region receives an average rainfall 1610 mm annually. The temperature reaches up to 40 degrees C in the months of May to June whereas, the higher peaks of the area receive frequent snowfall during winter season.

Vegetation Sampling and Analysis

The phytosociological analysis of the forests of study area (Yamuna valley) was carried out during June to August 2004 by using 10 x 10 m quadrats for trees. Each quadrat was subdivided in to 5 x 5 m sample plot for recording shrubs+saplings and 1 x 1 m for herbs+seedlings. The quadrats were laid out randomly throughout the study area. The size and the number of quadrats were determined by the species curve (Misra 1968) and the running means methods (Kershaw

1973). Thirty quadrats were randomly placed in the entire area, representing all the vegetation type and localities. In each quadrat, trees were recorded with >31.5 cm cbh (circumference at breast height i.e., 1.37 m above the ground) individually measured. Individuals within the cbh range of 10.5 to 31.4 cm were considered as shrubs+saplings and individuals < 10.5 cm cbh were considered as herbs+seedlings. The vegetation data were quantitatively analysed for abundance, density and frequency according to the formulae given by Curtis and Mc Intosh (1950) and Mishra (1968). The relative values were summed up to represent Importance Value Index (IVI) as per Curtis (1959). The distribution pattern of species was studied using the ratio of abundance to frequency if below 0.025 indicates regular distribution, between 0.025 - 0.050 indicates random distribution and when exceeds 0.050 indicates contagious distribution (Whitford 1949). The diversity index (H) was computed by using Shannon-Wiener information Index (Shannon and Wiener 1963). The concentration of dominance (CD) was computed by Simpson's Index (Simpson 1949). The dominance-diversity curves for trees, shrubs+saplings, and herbs+seedlings were drawn on the basis of importance value index (IVI). The utility index of important species of the sites were collected with concerned literature as well as the information collected from villagers.

Results and Discussion

Dominance

In the site-I of tree layer, a total of 33 species were observed. The dominant and co-dominant species were *Lannea coromandelica* and *Terminalia bellirica*, showing their values of IVI of 39.80 and 30.90 respectively, whereas, the highest (117 plants / ha) value of density was also recorded for *Lannea coromandelica*. Most of the species on the site showed contagious distribution pattern, except *Lannea coromandelica* which was randomly distributed (Table 1).

Species	Site-I			Site-II		
	Density	IVI	A/F	Density	IVI	A/F ratio
	(plant ha ⁻¹)		ratio	(plant ha ⁻¹)		
<i>Lannea coromandelica</i>	117	39.80	0.04	67	23.60	0.03
<i>Terminalia bellirica</i>	87	30.9	0.05	90	29.0	0.03
<i>Mallotus philippensis</i>	83	29.1	0.06	67	20.20	0.12
<i>Anogeissus latifolia</i>	40	17.0	0.07	100	29.5	0.05
<i>Ougeinia oojinensis</i>	33	14.6	0.12	7	4.36	0.15
<i>Toona serrata</i>	27	13.0	0.07	-	-	-
<i>Albizia lebbek</i>	27	12.6	0.15	10	5.69	0.23
<i>Boehmeria rugulosa</i>	20	11.80	0.45	3	3.05	0.30
<i>Litsea chinensis</i>	17	10.18	0.38	10	5.67	0.10
<i>Terminalia chebula</i>	17	9.63	0.06	33	13.6	0.05
<i>Terminalia alata</i>	17	9.26	0.17	23	9.67	0.13
<i>Moringa oleifera</i>	10	6.85	0.23	3	3.05	0.30
<i>Sapium insigne</i>	10	6.85	0.23	17	8.29	0.06
<i>Cordia vestita</i>	10	6.66	0.10	-	-	-
<i>Emblica officinalis</i>	10	6.66	0.10	7	4.36	0.15
<i>Schleichera oleosa</i>	7	6.48	0.06	-	-	-
<i>Bauhinia variegata</i>	7	5.18	0.15	10	5.69	0.23
<i>Ficus semicordata</i>	7	5.18	0.15	3	3.05	0.30
<i>Toona ciliata</i>	7	5.18	0.15	37	14.0	0.37
<i>Wrightia arborea</i>	7	5.18	0.15	3	3.04	0.30
<i>Bombax ceiba</i>	3	3.70	0.30	3	3.05	0.30
<i>Casearia graveolens</i>	3	3.70	0.30	3	3.05	0.30

Table 1. Phytosociological analysis of tree species on site-I and site-II

<i>Cassine glauca</i>	3	3.70	0.30	13	6.71	0.13
<i>Engelhardtia spicata</i>	3	3.70	0.30	-	-	-
<i>Erythrina suberosa</i>	3	3.70	0.30	-	-	-
<i>Ficus auriculata</i>	3	3.70	0.30	-	-	-
<i>Flacourtia indica</i>	3	3.70	0.30	-	-	-
<i>Holoptelea integrifolia</i>	3	3.70	0.30	3	3.05	0.30
<i>Jacaranda mimosifolia</i>	3	3.70	0.30	-	-	-
<i>Madhuca latifolia</i>	3	3.70	0.30	3	3.05	0.30
<i>Pinus roxburghii</i>	3	3.7	0.30	-	-	-
<i>Sterculia villosa</i>	3	3.7	0.30	-	-	-
<i>Adina cordifolia</i>	3	3.52	0.08	7	4.36	0.15
<i>Acacia catechu</i>	-	-	-	80	23.7	0.29
<i>Shorea robusta</i>	-	-	-	30	20.60	2.70
<i>Dalbergia sissoo</i>	-	-	-	33	13.0	0.33
<i>Toona serrata</i>	-	-	-	10	5.69	0.23
<i>Sterculia vilosa</i>	-	-	-	10	5.67	0.10
<i>Syzygium cumini</i>	-	-	-	10	5.67	0.10
<i>Cassia fistula</i>	-	-	-	7	4.36	0.15
<i>Ficus locar</i>	-	-	-	3	3.05	0.30
<i>Ficus religiosa</i>	-	-	-	3	3.05	0.30
<i>Legerstromia parviflora</i>	-	-	-	3	3.05	0.03
<i>Premna barbata</i>	-	-	-	3	3.05	0.30
Total	599	300.01		714	299.99	

Table 1. Phytosociological analysis of tree species on site-I and site-II contd.

In the tree layer of site-II a total of 34 tree species were recorded. Amongst the trees the highest value of density (100 plant ha⁻¹) and IVI (29.50) was recorded for *Anogeissus latifolia*. The co-dominant species of the site was *Terminalia bellirica* which showed their values of IVI (29.0) and density (90 plants / ha). The distribution pattern of all the species on the site was contagiously distributed except *Lannea coromandelica* and *Terminalia bellerica*, whose indicated random pattern of distribution.

In the shrub layer of site-I the highest value of IVI (26.70) was recorded for *Adhatoda vasica*, whereas, lowest (IVI-1.77) for *Agaveamericana*. In sapling layer the highest (140 plants / ha) value of density was recorded for *Boehmeria rugulosa* and the lowest for *Pyrus pashia* 2.0 plant / ha (Table 2).

Species	Site-I			Site-II		
	Density (plant ha ⁻¹)	IVI	A/F ratio	Density (plant ha ⁻¹)	IVI	A/F ratio
<i>Murraya koenigii</i>	233	27.6	0.11	163	22.79	0.06
<i>Adhatoda vasica</i>	230	26.7	0.14	190	24.80	0.07
<i>Woodfordia fruticosa</i>	173	22.5	0.09	163	21.99	0.08
<i>Colebrookia oppositifolia</i>	177	22.4	0.11	153	20.49	0.1
<i>Spermadictyon suaveolens</i>	160	20.7	0.12	13	18.6	0.1
<i>Boehmeria rugulosa</i> (Sapling)	140	18.3	0.2	-	-	-
<i>Rhamnus virgatus</i>	47	13.81	0.02	83	13.9	0.06
<i>Carissa spinarum</i>	77	12.0	0.19	97	14.49	0.11
<i>Lantana camara</i>	3	10.40	2.7	63	10.79	0.12
<i>Rhamnus procumbens</i>	3	10.40	2.7	27	5.89	0.27
<i>Desmodium cephalotes</i>	43	9.67	0.98	23	8.39	2.1
<i>Caesalpinia pectinata</i>	47	8.9	0.12	13	3.85	0.3
<i>Mallotus philippensis</i> (Sapling)	47	8.63	0.17	3	6.41	0.3
<i>Indigofera hexendrum</i>	47	8.56	0.26	-	-	-
<i>Rhus parviflora</i>	43	8.43	0.43	-	-	-
<i>Berberis asiatica</i>	27	5.98	0.27	3	1.71	0.3
<i>Ziziphus mauritiana</i>	23	5.84	0.53	10	3.18	0.23
<i>Pyrus pashia</i> (Sapling)	2	5.09	0.2	3	1.71	0.3
<i>Euphorbia royleana</i>	13	4.91	0.05	20	5.76	0.05
<i>Caseara graveolens</i> (Sapling)	10	3.93	0.9	13	5.05	1.2
<i>Ficus semicordata</i> (Sapling)	10	3.93	0.9	3	1.71	0.3

Table 2. Phytosociological analysis of shrub+sapling species on site-1 and site-II

<i>Nyctanthes arbor-tristis</i>	10	3.93	0.9	10	3.94	0.9
<i>Toona ciliata</i> (Sapling)	10	3.93	0.9	3	1.71	0.3
<i>Adina cordifolia</i> (Sapling)	7	2.85	0.6	-	-	-
<i>Bauhinia variegata</i> (Sapling)	7	2.85	0.6	7	2.83	0.6
<i>Cassia fistula</i> (Sapling)	7	2.85	0.6	7	2.83	0.6
<i>Emblica officinalis</i> (Sapling)	7	2.85	0.6	-	-	-
<i>Pogostemon benghalensis</i>	7	2.85	0.6	-	-	-
<i>Premna barbata</i> (Sapling)	7	2.85	0.6	-	-	-
<i>Toona serrata</i> (Sapling)	7	2.85	0.6	-	-	-
<i>Debregeasia latifolium</i>	7	2.65	0.15	13	3.84	0.13
<i>Millttia extensa</i> (Sapling)	3	2.01	0.08	7	2.83	0.6
<i>Agave americana</i>	3	1.77	0.3	-	-	-
<i>Ficus auriculata</i> (Sapling)	3	1.77	0.3	-	-	-
<i>Ficus palamata</i> (Sapling)	3	1.77	0.3	-	-	-
<i>Ougeinia oojeinensis</i> (Sapling)	3	1.77	0.3	7	2.83	0.6
<i>Sterculia villosa</i> (Sapling)	3	1.77	0.3	13	3.85	0.3
<i>Boehmeria platyphylla</i>	-	-	-	67	11.19	0.38
<i>Anogeissus latifolia</i> (Sapling)	-	-	-	33	7.82	0.75
<i>Eupatorium odoratum</i>	-	-	-	33	6.91	0.12
<i>Bauhinia vahlii</i> (Sapling)	-	-	-	30	6.52	0.11
<i>Artemisia vulgaris</i>	-	-	-	17	6.17	1.5
<i>Lannea coromandelica</i> (Sapling)	-	-	-	23	5.38	0.23

Table 2. Phytosociological analysis of shrub+sapling species on site-1 and site-II contd.

<i>Acacia catechu</i> (Sapling)	-	-	-	20	4.87	0.2
<i>Rubus ellipticus</i>	-	-	-	10	3.94	0.9
<i>Rubus nivens</i>	-	-	-	10	3.94	0.9
<i>Callicarpa macrophylla</i>	-	-	-	13	3.85	0.3
<i>Terminalia bellirica</i> (Sapling)	-	-	-	13	3.84	0.13
<i>Ficus locar</i> (Sapling)	-	-	-	7	2.89	0.6
<i>Cocculus laurifolius</i>	-	-	-	7	2.83	0.6
<i>Urtica dioica</i>	-	-	-	7	2.83	0.6
<i>Vitex negundo</i>	-	-	-	7	2.83	0.6
<i>Ficus religiosa</i> (Sapling)	-	-	-	3	1.71	0.3
<i>Sapium insigne</i> (Sapling)	-	-	-	137	1.71	0.3
<i>Syzygium cumini</i> (Sapling)	-	-	-	3	1.71	0.3
<i>Terminalia chebula</i> (Sapling)	-	-	-	3	1.71	0.3
<i>Erythrina suberosa</i> (Sapling)	-	-	-	3	1.17	0.3
Total	1649	300		1546	299.99	

Table 2. Phytosociological analysis of shrub+sapling species on site-1 and site-II contd.

In the shrub layer of site-II, a total of 25 species were recorded. Amongst the shrub species *Adhatoda vasica* showed highest value of density (190 plants / ha) and IVI (24.80). The lowest value of density (3.0 plants / ha) and IVI (1.71) was recorded for *Berberis aristata*. For the sapling layer, the maximum value of IVI was recorded for *Anogeissus latifolia* (7.82) and minimum (1.17) was recorded for *Erythrina suberosa* (Table 2)

In the herb layer on both the sites-I and II, the most dominant species was *Chrysopogen fulvus* while the least dominant species on site-I was *Artemisia vulgaris* and *Thalicturm foliolosum* and on site-II *Artemisia vulgaris* (Table 3). For seedling, in site-I only two seedlings (*Lansea coromandelica* and *Ougeinia oojenensis*) were recorded whereas, in site-II the total number of seedling were eight (Table 3).

Species	Site-I			Site-II		
	Density	IVI	A/F	Density	IVI	A/F
	(plant ha ⁻¹)		ratio	(plant ha ⁻¹)		ratio
<i>Chrysopogon fulvus</i>	373	27.70	0.31	217	21.08	0.16
<i>Dioscorea bulbifera</i>	160	18.0	0.07	137	19.00	0.04
<i>Euphorbia hirta</i>	200	18.0	0.18	133	14.6	0.19
<i>Achyranthes aspera</i>	160	15.90	0.14	57	8.58	0.1
<i>Nepeta elliptica</i>	140	13.5	0.26	47	8.08	1.05
<i>Desmodium cephalotes</i>	127	12.60	0.71	50	7.31	0.28
<i>Chrysopogon aciculatus</i>	117	12.10	0.21	50	7.7	0.13
<i>Parthenium hysterophorus</i>	117	11.7	0.42	230	22.1	0.14
<i>Apluda mutica</i>	97	10.30	0.54	-	-	-
<i>Aster thomsonii</i>	83	9.57	0.83	10	3.07	0.9
<i>Poa annua</i>	80	9.27	0.80	270	25.3	0.12
<i>Organum vulgare</i>	83	9.25	0.47	73	10.28	0.73
<i>Pogostemon benghalensis</i>	33	9.23	0.75	30	5.67	0.11
<i>Themeda arundinacea</i>	7	8.48	0.41	-	-	-
<i>Cassia tora</i>	60	8.27	1.35	60	10.0	1.35
<i>Euphorbia prostrata</i>	40	8.25	3.6	130	14.42	0.24
<i>Saccharum spontaneum</i>	63	7.96	0.23	90	11.8	0.1
<i>Reinwardtia indica</i>	53	7.51	1.2	-	-	-
<i>Tridax procumbens</i>	53	6.9	0.53	-	-	-
<i>Boehmeria platyphylla</i>	30	6.37	2.7	10	3.07	0.9
<i>Abrus precatorius</i>	40	5.99	0.9	-	-	-
<i>Viola serpens</i>	37	5.61	0.83	7	2.2	0.15

Table 3. Phytosociological analysis of herb+seedling species in site-I and site-II

<i>Agerattis houstoninum</i>	37	5.41	0.37	-	-	-
<i>Malva parviflora</i>	37	5.41	0.37	10	3.07	0.9
<i>Arisaema flavum</i>	27	4.90	0.15	37	7.06	0.07
<i>Thysanolaena maxima</i>	20	4.48	1.8	-	-	-
<i>Galainsonga parviflora</i>	27	4.47	0.6	-	-	-
<i>Euphorbia binata</i>	27	4.40	0.6	-	-	-
<i>Anemone sp.</i>	23	4.09	0.53	-	-	-
<i>Oxalis corniculata</i>	23	4.09	0.53	50	7.42	0.18
<i>Roscoea capitata</i>	23	4.09	0.53	-	-	-
<i>Cassia occidentalis</i>	17	3.85	1.5	17	4.71	1.5
<i>Salvia lanata</i>	10	2.60	0.9	-	-	-
<i>Staphania glabra</i>	10	2.60	0.9	30	5.36	0.17
<i>Artemisia vulgaris</i>	7	1.97	0.6	-	-	-
<i>Lansea coromendalica</i> (Seedling)	7	1.97	0.6	3	1.43	0.3
<i>Thalictrum foliolosum</i>	7	1.97	0.6	-	-	-
<i>Ougeinia oojeinensis</i> (Seedling)	3	1.34	0.3	3	1.43	0.3
<i>Clematis goriana</i>	-	-	-	30	7.99	2.7
<i>Phyllanthus armaris</i>	-	-	-	30	7.99	2.7
<i>Cyperus niveus</i>	-	-	-	53	7.91	0.53
<i>Cynodon dactylon</i>	-	-	-	53	7.63	0.3
<i>Ipomoea nil</i>	-	-	-	40	6.55	0.14
<i>Cannabis sativa</i>	-	-	-	20	5.53	1.8
<i>Solanum nigrum</i>	-	-	-	27	4.87	0.27
<i>Rumex hastatus</i>	-	-	-	17	3.67	0.38
<i>Cymbopogon jwarancus</i>	-	-	-	10	3.07	0.9
<i>Ficus religiosa</i> (Seedling)	-	-	-	10	3.07	0.9

Table 3. Phytosociological analysis of herb+seedling species in site-I and site-II contd.

<i>Urtica dioica</i>	-	-	-	10	2.69	0.23
<i>Ficus locar</i> (Seedling)	-	-	-	7	2.25	0.6
<i>Holarrhena pubescens</i> (Seedling)	-	-	-	7	2.25	0.6
<i>Ziziphus mauritiana</i>	-	-	-	7	2.18	0.60
<i>Woodfordia fruticosa</i>	-	-	-	7	2.05	0.60
<i>Acacia catechu</i> (Seedling)	-	-	-	3	1.43	0.3
<i>Anogeisuss latifolia</i> (Seedling)	-	-	-	3	1.43	0.3
<i>Milletia extensa</i> (Seedling)	-	-	-	3	1.43	0.3
Total	2458	300.01		2088	300	

Table 3. Phytosociological analysis of herb+seedling species in site-I and site-II contd.

Species diversity (H) and concentration of dominance (CD")

The diversity of study sites have been described in Table 4. The maximum (4.643) value of diversity for tree layer was recorded on site-II, whereas, minimum (4.580) was observed for site-I. In the shrub+sapling layer the maximum (5.021) and minimum (4.695) values of diversity were recorded on site-II and site-I respectively. In the herbs+seedlings layer, the minimum and maximum values of diversity were recorded on site-I and site-II respectively.

The lowest (0.053) and the highest (0.114) values of CD for tree layer was recorded on site-II and site-I. The ranged values for shrub+sapling and herb+seedling species were 0.040 to 0.049.

Dominance-diversity curve

Dominance-diversity curves plotted between importance value index and species sequences of trees, shrubs+saplings, herbs+seedlings (Figure 1-3) indicate a relationship between different species showing importance value on both sites.




Figure 1. Dominance-diversity curves for trees




Figure 2. Dominance-diversity curves for shrubs and saplings




Figure 3. Dominance-diversity curves for herbs and seedlings

In the present study two sites i.e. site-I and site-II were recorded because foothill belt of Garhwal Hiamalaya consist of thickly populated village and each village exploits forest resources for their basic needs from these forests (Kumar et al 2004). Due to over-exploitation and illicit felling as well as ruthless exploitation, most of the valuable species are either disappearing or rarely available.

The total density on site-I and site-II for trees were recorded 599 plants / ha and 714 plants / ha respectively. These values were quite lower than the values reported by Kumar et al (2004) to sub-tropical forests of Garhwal Himalaya. Some other tropical forests range of density 656 to 888 plants / ha was also reported by Kumar et al (2004). Several other workers also reported similar values for different forests of Garhwal Hiamalaya (Rajwar and Gupta 1992; Mishra et al. 2002)

The abundance to frequency ratio indicated that most of the species of shrub+sapling, herb+seedling and trees were contagiously distributed except few species of trees, which showed random distribution pattern. Contagious distribution in natural vegetation has been reported by Greigh-Smith (1957), Kershaw (1973). Similar finding also reported by Kumar et al (2004) for tropical forests of Garhwal Himalaya.

In present study the range of diversity were 4.580 to 4.643 (for trees), 4.695 to 5.021 (for shrubs+saplings), and 4.962 to 4.986 (for herbs+seedlings). The ranges of diversity were high on the sites due to disturbance and invasion by new species. In Table 4 indicates that increasing diversity and reduced concentration of dominance has been shown to be associated with increased stability (Mc Naughton 1967). Pandey and Singh (1985) have also reported increasing species diversity in disturbed ecosystem of Kumaon Himalaya. Disturbance has always been considered causes invasion and species replacement, altered, growth, establishment, germination etc. the disturbance affects the stability of the ecosystem and the successional process of the area. Disturbance caused by sever grazing arrests the ecological determinations. However, grazing animals which consume the ground vegetation plays positive role in dispersal, establishment and growth of such species by scarification and pelletization of seed (with dung).

The dominance diversity (d-d) curves (based on IVI) approach a geometric series for all the strata. Mostly these curves follow the geometric series in conformity with niche pre-emption hypothesis (Motonura 1934). The geometric form in often shown by vascular plants having lower density (Whittaker 1975)

Uses and Conservation Methods

Forest is a living resource. A large number people of the hill and plain area depends on forest resources. Forest constitutes the richest resources amongst the other resources of the world. They are the vital component to sustaining the life support system on earth. Forests have been playing a pivotal role in the socio-economic development of a country or region. They are the important source of subsistence, employment, revenue earning and raw material to various Industrial uses. There role in ecological balance, environmental stability, biodiversity conservation, food security and sustainable development have been widely recognized. Forests are the source of various components, e.g., food, fodder, fiber, medicinal, tannin and oils, gum and many other things for human benefits. Generally it is observed that the forest areas in the vicinity of the villages have been degrading much faster rate than the forests growing far from the village locality. It is due to easily accessible to the villagers for their basic needs. As per the data collected from local inhabitants and concern literature revels that the forest resources of the area is rich for human benefits (Table 5). Therefore, there is an urgent need to conserve the forest resource both macro and molecular level for human and sustainable development of environment, besides that awareness to the villagers is essential how they can develop suitable techniques for sustainable utilization of forest resources.

Table 4. Species diversity, concentration of dominance of different layers on site-I and site-II

Sites/Layers	Species diversity	Concentration of dominance
Site-1		
<i>Tree</i>	4.580	0.114
<i>Sapling+shrub</i>	4.695	0.049
<i>Seedling+herb</i>	4.962	0.039
Site-II		
<i>Tree</i>	4.643	0.053
<i>Sapling+shrub</i>	5.021	0.040
<i>Seedling+herb</i>	4.986	0.039

Table 5. Utility Index of plant species recorded from the site-I and site-II

Species	Utility Index
<i>Acacia catechu</i>	Medicinal, Fodder, Dye, Gum, Lac, Kath formation
<i>Achyranthes aspera</i>	Local beverage, Medicinal.
<i>Adhatoda vasica</i>	Young leaves for cough and cold
<i>Adina cordifolia</i>	Toy and handicraft.
<i>Albizia lebbeck</i>	Fodder, Paper and pulp, Lac, Medicinal
<i>Anogeissus latifolia</i>	Fodder, Fiber
<i>Apluda mutica</i>	Thatching huts and brooms
<i>Artemisia vulgaris</i>	Medicinal (leaf juice use intestinal problem)
<i>Bauhinia variegata</i>	Fodder Dye, Ornamental, Oil and fat
<i>Berberis asiatica</i>	Medicinal plant, Fodder
<i>Bombax ceiba</i>	Fodder Fiber, Forage Ornamental, Oil and fat
<i>Carissa spinarum</i>	Commercial, Tanning
<i>Cassia fistula</i>	Dye, Ornamental, Medicinal
<i>Colebrookia oppositifolia</i>	Leaves used as medicinally
<i>Cynodon dactylon</i>	Plant used un several religious ceremonies
<i>Dalbergia sissoo</i>	Fodder, Toy and handicraft, Paper and pulp, Resin used in skin ailments
<i>Embllica officinalis</i>	Fodder, Dye, Oil and fat, Source of Vitamin and ingredient of Trifala
<i>Erythrina suberosa.</i>	Fodder, Fiber

<i>Ficus auriculata</i>	Fodder Fiber
<i>Ficus religiosa</i>	Fodder, Fiber, Commercial, Lac, Religious plant
<i>Holoptelea integrifolia</i>	Fodder, Bark used as rheumatic pain
<i>Jacaranda mimosifolia</i>	Ornamental purpose
<i>Lannea coromandelica</i>	Fiber. Dye, Gum
<i>Lantana camara</i>	Leaves insecticidal or germicidal
<i>Madhuca longifolia</i>	Fodder, Medicinal
<i>Mallotus philippensis</i>	Dye, Fruit, Fodder, Medicinal
<i>Moringa oleifera</i>	Fodder, Fiber Oil and fat, Fruit edible, Medicinal
<i>Murraya koenigii</i>	Fodder, Toy and handicraft, bark, root and leaves used medicinally
<i>Nyctanthes arbor-tristis</i>	Fodder, Medicinal, Dye
<i>Organum vulgare</i>	Medicinal (carminative, diuretic , diaphoretic)
<i>Ougeinia oojenensis</i>	Fodder, Gum used as digestive troubles
<i>Pinus roxburghii</i>	Fodder, Dye
<i>Rhus parviflora</i>	Fodder, Medicine (cholera)
<i>Schleichera oleosa</i>	Fodder Lac, Oil and fat Medicinal
<i>Shorea robusta</i>	Fodder, Commercial, Dye, Resin, Silk
<i>Syzygium cumini</i>	Fodder, Dye, Silk
<i>Terminalia chebula</i>	Medicinal, fodder, dye.
<i>Terminalia alata</i>	Dye, Silk, Paper and pulp
<i>Terminalia bellirica</i>	Medicinal, Fodder, Dye, Resin, Silk
<i>Toona ciliata</i>	Fodder, Fuel Oil and Medicinal
<i>Vitex negundo</i>	Medicinal (roots, leaves, fruits)
<i>Woodfordia fruticosa</i>	Fodder, Medicinal, dried flower as tonic
<i>Ziziphus mauritiana</i>	Fodder, Fruit edible

The methods basically used for conservation of biological diversity both (animals and plants) are *in-situ* and *ex-situ* conservation. *Ex-situ* conservation of species done outside the area where they are naturally growing (e.g., botanical, horticulture and recreational gardens) while, *in-situ* conservation is done in the natural habitats (e.g., area demarcated by under rules are National Parks, Sanctuaries, Biosphere Reserve, Preservation plots etc.). Besides this several other traditional methods are used by peoples can also considered i.e., (1). Sacred groves forest should be declared (2) Large scale cultivation of economic and medicinal valuable species should be encouraged to local inhabitants (3) Public awareness (4) Encouragement of social forestry programme for some essential species, for fuel, fodder and other purposes (5) Deforestation must be stopped by adopting law and regulation, should be developed by local village community and government level. (6) Plantation

programmes should be made as compulsory for all villagers of the area.

Keeping all the consideration for sustainable development of forest resources, emphasize should be given to develop suitable agroforestry, models specially selected multipurpose tree species in conjunction with agricultural crops over the same unit of land to maximize productivity and sustainable utilization of these resources.

References

- Bargali, K., Usman and. Joshi, M. 1998. Effect of forest covers on certain site and soil characteristics in Kumaun Himalayas. *Indian Journal of Forestry*, 21 (3): 224-227.
- Canham, C.D. and Marks, P. L. 1985. The response of woody plants to disturbance patterns of establishment and growth. In: *The Ecology of Natural Disturbance and Patch Dynamics* (eds. S.T.A. Pickett and P.S. White), Academic Press, Orlando, Fla..
- Curtis, J.T. 1959. *The Vegetation of Wisconsin*. An Ordination of plant communities, University Wisconsin press, Madison Wisconsin, 657pp.
- Curtis, J.T. and mc Intosh, R. P. 1950. The Interrelation of certain analytic and synthetic phytosociological characters. *Ecology* 31: 434-455.
- Gaur, R.D. 1982. Dynamics of vegetation of Garhwal Himalayas. *The Vegetational Wealth of the Himalayas* (ed. G.S. Paliwal), pp. 12-25.
- Greig-Smith, P. 1957. *Quantitative Plant Ecology*, 2nd edition. Butterworth, London.
- Kershaw, K.A. 1973. *Quantitative and Dynamic Plant Ecology* (2nd Edition), ELBSD& Edward Arnold, London.
- Kumar, M., Sharma C.M. and Rajwar, G.S. 2004. A study on the community structure and diversity of a sub-tropical forest of Garhwal Himalayas. *Indian Forester* 130 (2): 207-214
- Loucks, O. L., Ek, A. R., Johanson, W. C. and Monserud R. A. 1980. Growth ageing and succession. In: D. E. Reichle (Editor). *Dynamics properties of Forest Ecosystem*. International Biological programme 23, Combridge University Press, Malta, pp. 37-85.
- Mc Naughton, S. J. 1967. Relationship among functional properties of California grassland. *Nature* 216: 168-169.
- Mishra, A., Mishra, L. Sharma, C.M. and Sharma, S.D. 2002. Community structure and growing stock variation on different aspects of dry semi-evergreen *Shorea robusta* forest cover type of Garhwal Himalaya. *Annals of forestry* 10 (2): 233-242.
- Mishra, R.1968. *Ecology Workbook*. Oxford & IBH, Calcutta.
- Motomura, I.1932. A statistical treatment of associations (in Japanese). *Japanese Journal of Ecology* 44: 379-383.
- Osmaston, A.E. 1922 Notes on the forest communities of the Garhwal Himalaya. *Journal of Ecology* 10: 129-187.
- Pandey, A.N. and Singh, J.S. 1985. Mechanism of ecosystem recovery: A case study of Kumaun Himalaya. *Recreation and Revegetation Research* 3: 271-292
- Pickett, S. T. A. and White, P. S. 1985: *The Ecology of Natural Disturbance and Patch Dynamics*. Academic Press, Orlando, Fla.
- Puri, G.S. 1960. *Indian Forest Ecology* 2 Vol. New Delhi.
- Rajwar, G.S. and Gupta S.K. 1992. Structure of forest vegetation of Garhwal Siwalik Hills between Khoh and Ganga. *Indian Forester* 118 (2): 148-165.
- Ralhan, P.K., Saxena, A. K. and Singh, J. S. 1982. Analysis of forest vegetation at and around Nainital in Kumaun Himalaya. *Proc. Indian National Sciences Academy* 348:121-137.
- Saxena, A. K., Singh, S.P. and Singh, J.S. 1984. Population structure of forests of Kumaun Himalaya: implications for management. *Journal of Environmental Management* 19: 307-324
- Saxena, A.K. and Singh, J.S. 1982. A phytosociological analysis of woody species in forest communities of a part of Kumaun Himalaya. *Vegetatio* 50: 3-22 .
- Saxena, A.K. Singh, J.S. 1982.A phytosociological analysis of woody species in forest communities of a part of Kumaun Himalaya. *Vegetatio* 50: 3-22
- Saxena, A.K. Singh, S.P. and Singh, J.S. 1984. Population structure of forests of Kumaun Himalaya: Implications for Management. *Journal of Environment Management* 19: 307-324.
- Shannon, C.E. and Wiener, W. 1963. *The Mathematical Theory of Communication*. University of Illinois press, Urbana. 117 pp.
- Simpson, E. H. 1949. *The Measurement of Diversity*. *Nature*. 163:688.
- Singh, J.S., Singh, S.P. and Shastri, C. 1981. *Science of rural development in mountains*.

Gyanodaya Prakashan, Nainital, India.

Whitemore, T.C., 1984. Tropical rain forest dynamics and its implications for management. In: *Rain Forest Regeneration and Management*. (Ed. A. Gomez-pompa, T.C. Whitemore and M. Hadley), Man Biosphere Series 6

Whitford, P.B. 1949. Distribution of woodland plants in relation to succession and clonal growth. *Ecology* 30: 199-208.

Whittaker, R.H. 1975. *Communities and Ecosystems*, 2nd edition. Macmillan Publishing Co., New York. 385.